


WHAT IS CLAIMED IS:

1. A compound having the formula:



wherein R¹ is lower alkyl, bridged alkyl, aryl, heteroaryl, aralkyl, cycloalkyl, a 5- or 6-membered saturated heterocycle, C₁₋₄alkylenearyl, C₁₋₄alkyleneOaryl, C₁₋₄alkyleneheteroaryl, C₁₋₄alkylene-Het, C₂₋₄alkylenearyloaryl, C₁₋₄alkylene bridged alkyl, C₁₋₃alkylenecycloalkyl, substituted or unsubstituted propargyl, substituted or unsubstituted allyl, or halocycloalkyl;

R² is hydrogen, methyl, or halo-substituted methyl;

R³ is selected from the group consisting of C(=O)OR⁷, C(=O)R⁷, C(=NH)NR⁸R⁹, C(=O)NR⁸R⁹, lower alkyl, bridged alkyl, cycloalkyl, haloalkyl, halocycloalkyl, C₁₋₃alkylenecycloalkyl, a 5- or 6-membered saturated heterocycle, aryl, heteroaryl, C₁₋₃alkyleneC(=O)R⁷, C(=O)C(=O)NR⁸R⁹, C₁₋₄alkyleneOR⁷, C₁₋₃alkylenearyl, SO₂heteroaryl, Het, aralkyl, alkaryl, heteroaralkyl, heteroalkaryl, C₁₋₃alkyleneC(=O)OR⁷, C(=O)C₁₋₃alkyleneC(=O)OR⁷, C₁₋₃alkyleneheteroaryl, C(=O)C(=O)OR⁷, C(=O)C₁₋₃alkyleneC(=O)OR⁷, C(=O)-

C₁₋₃alkyleneNH(C=O)OR⁷, C(=O)C₁₋₃alkyleneNH₂, and
NHC(=O)OR⁷;

R⁴ is hydrogen, lower alkyl, haloalkyl,
cycloalkyl, or aryl;

R⁵ is hydrogen, lower alkyl, alkynyl, halo-
alkyl, cycloalkyl, or aryl;

R⁶ and R¹⁰, independently, are hydrogen,
lower alkyl, aralkyl, SO₂R⁶, or C(=O)R⁷;

R⁷ is selected from the group consisting of
branched or unbranched lower alkyl, heteroaryl, a
heterocycle, aralkyl, and aryl, and R⁷ can be option-
ally substituted with one or more of RO⁸, NR⁸R⁹, or
SR⁸;

R⁸ and R⁹, same or different, are selected
from the group consisting of hydrogen, lower alkyl,
cycloalkyl, aryl, heteroaryl, alkaryl, heteroaralk-
yl, heteroalkaryl, and aralkyl, or R⁸ and R⁹ can be
taken together form a 4-membered to 7-membered ring;


R¹⁰ is hydrogen, alkyl, haloalkyl, cyclo-
alkyl, aryl, C(=O)alkyl, C(=O)cycloalkyl, C(=O)aryl,
C(=O)Oalkyl, C(=O)Ocycloalkyl, C(=O)aryl, CH₂OH,
CH₂Oalkyl, CHO, CN, NO₂, or SO₂R¹¹;

R¹¹ is alkyl, cycloalkyl, trifluoromethyl,
aryl, aralkyl, or NR⁸R⁹;

salts and solvates thereof.




20250154-024502






2. The compound of claim 1 having the structure:






20251201 15:22:00

3. The compound of claim 1 wherein R¹ is selected from the group consisting of:

























1077454-024502

- 110 -



20250215 15:02:54





- 113 -





H-

CH₃-







(CH₃)₃C-







(CH₃)₃C(CH₂)₂


(CH₃)₂C(CH₂)₂-
|
OCH₃




1007154-021502














and













10077154.021502











4. The composition of claim 1 wherein R¹ is selected from the group consisting of:







1007154-021502













10077154-021502

























1007154-021502










10077154-021502

















1077154-021502




1007154-021502





and



1077154-121502

5. The compound of claim 1 wherein R' is selected from the group consisting of hydrogen, methyl, trifluoromethyl, cyclopropyl, ethynyl, and phenyl.

6. The compound of claim 1 wherein R is hydrogen or lower alkyl.


7. The compound of claim 1 wherein R' is selected from the group consisting of hydrogen, $C(=O)R^7$, $C(=O)OR^7$, ethyl, benzyl, SO_2CH_3 , and $SO_2C_6H_5$.

8. The compound of claim 1 wherein R⁷ is lower alkyl.

9. The compound of claim 1 wherein R⁸ and R⁹, independently, are hydrogen or lower alkyl, or are taken together form a 5-membered or 6-membered ring.

10. The compound of claim 1 wherein R⁻² is selected from the group consisting of hydrogen and lower alkyl.

11. The compound of claim 1 wherein R¹ is selected from the group consisting of cyclopentyl, cyclopropylmethyl, tetrahydrofuryl, indanyl, norbornyl, phenethyl, and phenylbutyl; R² is selected from the group consisting of methyl and difluoromethyl; R³ is selected from the group consisting of benzyl, CO₂CH₃, C(=O)CH₂OH, C(=O)CH(CH₃)OH, C(=O)C(CH₃)₂OH, and



R⁴ is hydrogen; R⁵ is hydrogen or methyl; R⁶ is selected from the group consisting of hydrogen, methyl, ethyl, benzoyl, SO₂CH₃, SO₂C₆H₅, benzyl, C(=O)C(CH₃)₃, and acetyl; R⁷ is hydrogen or methyl; R⁸ is methyl; and R⁹ is hydrogen.

1007154-021502

12. The compound of claim 1 selected from the group consisting of

Methyl (4S,3R)-4-(3-cyclopentyloxy-4-methoxyphenyl)-3-methyl-3-{[benzylamino]methyl}pyrrolidine carboxylate

Methyl (4S,3R)-3-(aminomethyl)-4-(3-cyclopentyloxy-4-methoxyphenyl)-3-methylpyrrolidinecarboxylate

Methyl (3S,4S)-4-(3-cyclopentyloxy-4-methoxyphenyl)-3-methyl-3-{[methylsulfonyl]amino}methylpyrrolidinecarboxylate

Methyl (4S,3R)-3-[(acetylamino)methyl]-4-(3-cyclopentyloxy-4-methoxyphenyl)-3-methylpyrrolidinecarboxylate

Methyl (4S,3R)-4-(3-cyclopentyloxy-4-methoxyphenyl)-3-methyl-3-[(phenylcarbonylamino)methyl]pyrrolidinecarboxylate

Methyl (3S,4S)-4-(3-cyclopentyloxy-4-methoxyphenyl)-3-methyl-3-{[phenylsulfonyl]amino}methylpyrrolidinecarboxylate

Bis{[(4S,3R)-4-(3-cyclopentyloxy-4-methoxyphenyl)-3-methyl-1-carboxymethylpyrrolidin-3-yl]methyl}amine

1-[(3S,4S)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-3-methyl-1-benzylpyrrolidin-3-yl]ethylamine

1007154 021502

1-[(3S,4S)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-3-methyl-1-benzylpyrrolidin-3-yl]ethylamine

N-{1-[(3S,4S)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-3-methyl-1-benzylpyrrolidin-3-yl]ethyl}benzamide

N-{1-[(3S,4S)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-3-methyl-1-benzylpyrrolidin-3-yl]ethyl}benzamide

N-{1-[(3S,4S)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-3-methyl-1-benzylpyrrolidin-3-yl]ethyl}acetamide

N-{1-[(3S,4S)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-3-methyl-1-benzylpyrrolidin-3-yl]ethyl}acetamide

3-(S)-(1-Acetylaminoethyl)-4-(S)-(3-cyclopentyloxy-4-methoxyphenyl)-3-methylpyrrolidine-1-carboxylic acid methyl ester

{1-[(3S,4S)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-3-methyl-1-benzylpyrrolidin-3-yl]ethyl}-(phenylsulfonyl)amine

{1-[(3S,4S)-4-(3-cyclopentyloxy-4-methoxyphenyl)-3-methyl-1-benzylpyrrolidin-3-yl]ethyl}-(phenylsufonyl)amine

{1-[(3S,4S)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-3-methyl-1-benzylpyrrolidin-3-yl]ethyl}-(methylsulfonyl)amine










10077154-021502

{1-[(3S,4S)-4-(3-Cyclopentyloxy-4-methoxyphenyl)-3-methyl-1-benzylpyrrolidin-3-yl]ethyl}-
(methylsulfonyl)amine, and

Methyl (3S,4S)-4-(3-cyclopentyloxy-4-methoxyphenyl)-
3-methyl-3-[(methylamino)ethylpyrrolidine carbox-
ylate.

1007154-021502

13. The compound of claim 1 selected from the group consisting of



14. The compound of claim 1 having an IC_{50} vs. human recombinant PDE4 of about 1 nM to about 25 μ M.

15. The compound of claim 1 having a PBL/TNF α EC_{50} of about 10 nM to about 20 μ M.

16. The compound of claim 1 having an IC_{50} vs. human recombinant PDE4 of about 1 nM to about 25 μ M, and a PBL/TNF α EC_{50} of about 10 nM to about 25 μ M.

17. The compound of claim 1 having an IC_{50} vs. human recombinant PDE4 of about 100×10^{-6} M or less.

18. The compound of claim 1 having an IC_{50} vs. human recombinant PDE4 of about 50×10^{-6} M or less.

19. The compound of claim 1 having a PBL/TNF α EC_{50} of about 5 μ M or less.

20. The compound of claim 1 having a PBL/TNF α EC_{50} of about 2 μ M or less.

21. The compound of claim 1 having an IC_{50} vs. human recombinant PDE4 of about 100×10^{-6} or less and a PBL/TNF α EC_{50} of about 5 μ M or less.

22. The compound of claim 1 having an IC_{50} vs. human recombinant PDE4 of about 50×10^{-6} or less and a PBL/TNF α EC_{50} of about 2 μ M or less.

23. A pharmaceutical composition comprising a compound of claim 1, a pharmaceutically acceptable carrier, and, optionally, a second antiinflammatory therapeutic agent.

24. The composition of claim 23 wherein the second antiinflammatory therapeutic agent is capable of targeting TNF α .

25. A method of treating a mammal having a condition where inhibition of a cAMP-specific PDE is of therapeutic benefit, said method comprising administering to said mammal at therapeutically effective amount of a compound of claim 1.

26. A method of modulating cAMP levels in a mammal comprising administering to said mammal an effective amount of a compound of claim 1.

27. A method of treating a mammal having a condition where inhibition of a cAMP-specific PDE is of a therapeutic benefit comprising administering to said mammal an effective amount of a pharmaceutical composition comprising a compound of claim 1 and a pharmaceutically acceptable carrier.

28. The method of claim 27 wherein the condition is an allergic disease, an autoimmune disease, an inflammatory disease, an arthritic disease, or dermatitis.

29. The method of claim 27 wherein the condition is rheumatoid arthritis, osteoarthritis, gouty arthritis, or spondylitis.

30. The method of claim 27 wherein the condition is thyroid-associated ophthalmopathy, Behcet disease, sepsis, septic shock, endotoxic shock, gram negative sepsis, gram positive sepsis, toxic shock syndrome, allergic conjunctivitis, vernal conjunctivitis, or eosinophilic granuloma.

31. The method of claim 27 wherein the condition is asthma, chronic bronchitis, allergic rhinitis, adult respiratory distress syndrome, chronic pulmonary inflammatory disease, chronic obstructive pulmonary disease, silicosis, or pulmonary sarcoidosis.

32. The method of claim 27 wherein the condition is reperfusion injury of the myocardium, brain or extremities as a brain or spinal cord injury due to trauma.

33. The method of claim 27 wherein the condition is a fibrosis, keloid formation, or scar tissue formation.

34. The method of claim 27 wherein the condition is systemic lupus erythematosus, a transplant rejection disorder, a graft vs. host reaction, or an allograft rejection.

2025 JUN 11 10 02 AM

35. The method of claim 27 wherein the condition is chronic glomerulonephritis, an inflammatory bowel disease, Crohn's disease, or ulcerative colitis.

36. The method of claim 27 wherein the condition is proliferative lymphocytic disease or a leukemia.

37. The method of claim 27 wherein the condition is an inflammatory dermatosis, atopic dermatitis, psoriasis, or urticaria.

38. The method of claim 27 wherein the condition is a cardiomyopathy, congestive heart failure, atherosclerosis, pyrexia, cachexia, cachexia secondary to infection or malignancy, cachexia secondary to acquired immune deficiency syndrome, ARC, cerebral malaria, osteoporosis, a bone resorption disease, fever and myalgias due to infection, erectile dysfunction, diabetes insipidus, a central nervous system disorder, depression, multi-infarct dementia, an anxiety or stress response, cerebral ischemia, tardive dyskinesia, Parkinson's disease, or premenstrual syndrome.

39. The method of claim 27 wherein the mammal exhibits a minimal emetic response.

40. The method of claim 27 wherein the mammal is free of an emetic response.

10077154-001502
2025120-4572001

41. The method of claim 27 wherein the mammal exhibits minimal adverse central nervous system side effects.

42. The method of claim 27 wherein the mammal is free of adverse central nervous system side effects.

43. The method of reducing TNF levels in a mammal comprising administering to said mammal therapeutically effective amount of a compound of claim 1.

44. A method of suppressing inflammatory cell activation in a mammal comprising administering to said mammal a therapeutically effective amount of a compound of claim 1.

45. A method of inhibiting PDE4 function in a mammal comprising administering to said mammal a therapeutically effective amount of a compound of claim 1.

20250215 02:50